



Are eccentricity fluctuations able to explain the centrality dependence of v_4 ?

Clément Gombeaud, Jean-Yves Ollitrault

► To cite this version:

Clément Gombeaud, Jean-Yves Ollitrault. Are eccentricity fluctuations able to explain the centrality dependence of v_4 ?. Journal of Physics G: Nuclear and Particle Physics, 2010, 37 (9), pp.94024. 10.1088/0954-3899/37/9/094024 . hal-00600843

HAL Id: hal-00600843

<https://hal.science/hal-00600843>

Submitted on 16 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Are eccentricity fluctuations able to explain the centrality dependence of v_4 ?

Clément Gombeaud and Jean-Yves Ollitrault

Institut de physique théorique CEA-Saclay 91191 Gif-sur-Yvette, France

E-mail: clement.gombeaud@cea.fr

Abstract.

The fourth harmonic of the azimuthal distribution of particles v_4 has been measured for Au-Au collisions at the Relativistic Heavy Ion Collider (RHIC). The centrality dependence of v_4 does not agree with the prediction from hydrodynamics. In particular, the ratio $v_4/(v_2)^2$, where v_2 denotes the second harmonic of the azimuthal distribution of particles, is significantly larger than predicted by hydrodynamics. We argue that this discrepancy is mostly due to elliptic flow (v_2) fluctuations. We evaluate these fluctuations on the basis of a Monte Carlo Glauber calculation. The effect of deviations from local thermal equilibrium is also studied, but appears to be only a small correction. Combining these two effects allows us to reproduce experimental data for peripheral and midcentral collisions. However, we are unable to explain the large magnitude of $v_4/(v_2)^2$ observed for the most central collisions.

1. Introduction

The azimuthal distribution of emitted particles is a good tool for understanding the bulk properties of the matter created in non central nucleus-nucleus collisions. In the center of mass rapidity region, it can be expanded in Fourier series:

$$\frac{dN}{d\phi} \propto 1 + 2v_2 \cos(2\phi) + 2v_4 \cos(4\phi) + \dots \quad (1)$$

where ϕ is the azimuthal angle with respect to the direction of the impact parameter, and odd harmonics are zero by symmetry. The large magnitude of elliptic flow v_2 observed at RHIC suggests that the matter created in Au-Au collisions behaves like an almost perfect fluid. However, recent experiments [1, 2] observed that, at midrapidity and fixed p_t , $v_4 \simeq (v_2)^2$, while ideal hydrodynamics predicts that $v_4 = \frac{1}{2}(v_2)^2$ [3]. In this contribution, I investigate this discrepancy.

2. Fluctuations in initial conditions




Figure 1. (Color online) Picture of the two frames used for defining the initial eccentricity (from [4]). The x axis defines the reaction plane while the x' axis is the minor axis of the ellipse drawn by the participating nucleons (grey dots).

Figure 1 presents a schematic picture of a non central heavy-ion collision (HIC). The overlap area of the nuclei has an almond shape, which generates elliptic flow. However, the matter is not continuously distributed in a nucleus. The positions of the nucleons in the colliding nucleus are important: they also draw an ellipse which differs from the overlap area both in eccentricity and in orientation. From one event to the other, even at fixed impact parameter, the positions of the nucleons in the nucleus fluctuate. The participant plane eccentricity (ϵ_{PP}), defined as the eccentricity of the ellipse drawn by the participating nucleons [5, 6], thus fluctuates. Since elliptic flow appears to be driven by this participant plane eccentricity, these eccentricity fluctuations translate into fluctuations of the flow coefficients v_2 and v_4 [7].

3. Modeling eccentricity fluctuations

The initial distribution of energy, which is needed to compute the initial eccentricity in a HIC, is poorly known. In this talk I use a specific model, based on a Monte Carlo Glauber (MCG) calculation [8]. The initial eccentricity is given for each event by:

$$\epsilon_{PP} = \frac{\sqrt{(\sigma_y^2 - \sigma_x^2)^2 + 4\sigma_{xy}^2}}{\sigma_x^2 + \sigma_y^2} \quad (2)$$

where $\sigma_x^2 = \langle x^2 \rangle - \langle x \rangle^2$ and $\sigma_{xy} = \langle xy \rangle - \langle x \rangle \langle y \rangle$ and the $\langle \rangle$ denote averages over participating nucleons. Each participant nucleon is given a weight proportional to the number of particles it creates, according to the two-component picture: $w = (1 - x) + xN_{\text{coll-nucleon}}$ where $N_{\text{coll-nucleon}}$ is the number of binary collisions of the nucleon. The sum of weights scales like the multiplicity:

$$\frac{dN_{ch}}{d\eta} = n_{pp} \left[(1 - x) \frac{N_{part}}{2} + xN_{coll} \right]. \quad (3)$$

where N_{part} and N_{coll} are respectively the number of participants and of binary collisions of the considered event. We choose the value $x = 0.13$ which best describes the charged hadron multiplicity observed experimentally [9]. We define the centrality according to the number of participants. We evaluate eccentricity fluctuations in centrality classes containing 5% of the total number of events. We do not introduce any hard core repulsion between nucleons in the MCG.

4. How eccentricity fluctuations affect v_4/v_2^2 .




Figure 2. (Color online) Centrality dependence of $v_4/(v_2)^2$: data taken from STAR [10] and PHENIX [11]; error bars on STAR data points are our estimates of non-flow errors [12]. Lines are predictions from ideal hydro with or without fluctuations.

There is no direct measure of the flow coefficients v_2 and v_4 . They can be obtained using different analysis methods. The one I will consider now relies on

azimuthal correlations between particles near midrapidity. Experimentally, v_2 can be extracted from the 2-particle correlation and v_4 from the 3-particle correlation using $\langle \cos(2\phi_1 - 2\phi_2) \rangle = \langle (v_2)^2 \rangle$ and $\langle \cos(4\phi_1 - 2\phi_2 - 2\phi_3) \rangle = \langle v_4(v_2)^2 \rangle$, where angular brackets denote an average value within a centrality class. Thus, any experimental measure of v_4/v_2^2 obtained using this method is rather a measure of $\langle v_4 \rangle / \langle v_2^2 \rangle^2$. Taking into account the ideal hydrodynamics prediction $v_4 = \frac{1}{2}(v_2)^2$ [3], we obtain:

$$\left(\frac{v_4}{(v_2)^2} \right)_{\text{measured}} = \frac{1}{2} \frac{\langle (v_2)^4 \rangle}{\langle (v_2)^2 \rangle^2} > \frac{1}{2}. \quad (4)$$

Assuming that v_2 scales like the participant plane eccentricity ϵ_{PP} , the effects of fluctuations on v_4/v_2^2 is obtained by computing:

$$\left(\frac{v_4}{(v_2)^2} \right)_{MCG} = \frac{1}{2} \frac{\langle \epsilon_{PP}^4 \rangle}{\langle \epsilon_{PP}^2 \rangle^2}. \quad (5)$$

The resulting prediction for v_4/v_2^2 is displayed in figure 2. Fluctuations clearly explain most of the difference between hydro and data. It also appears that experimental data are still slightly higher than our prediction from fluctuations. However, these predictions are based on a Monte-Carlo Glauber parametrization of the initial conditions, predictions may differ with another parametrization.

5. Flow fluctuations from experimental data




Figure 3. Same as figure 2, additional points (labeled "fluct. from v_2 ") are obtained using equation (6) with $v_2\{4\}$ from [13] and $v_2\{2\}$ from [14].

Another possible way of evaluating flow fluctuations is to compare the values of v_2 obtained using different analysis methods. Elliptic flow can be obtained from both 2-particle cumulants ($v_2\{2\}$) and from 4-particle cumulants ($v_2\{4\}$) using $v_2\{2\}^2 = \langle (v_2)^2 \rangle$

(neglecting the non-flow contribution) and $v_2\{4\}^4 = 2\langle(v_2)^2\rangle^2 - \langle(v_2)^4\rangle$. Inverting this equation leads to:

$$\left(\frac{v_4}{(v_2)^2}\right) = \frac{1}{2} \frac{\langle(v_2)^4\rangle}{\langle(v_2)^2\rangle^2} = \frac{1}{2} \left(2 - \left(\frac{v_2\{4\}}{v_2\{2}\}\right)^4\right). \quad (6)$$

The values of v_4/v_2^2 obtained using this method are displayed on figure 3. They overshoot slightly our results from MCG eccentricity fluctuations, but the overall agreement remains good. This provides a good check of our MCG prediction. A small residual discrepancy remains between our prediction and the experimental data. We argue that for peripheral to midcentral collisions, it may be understood in terms of deviations from local thermal equilibrium.

6. Partial thermalization effects

So far, I have only discussed how fluctuations in initial conditions modify the prediction from ideal hydrodynamics for v_4/v_2^2 . But ideal hydrodynamics relies on the very strong assumption that the system remains in local thermal equilibrium (a regime where the average number of collisions per particle n_{coll} is large) throughout the evolution. In a previous work [15] we have shown that, in order to reproduce the centrality dependence of elliptic flow, the deviation from local thermal equilibrium must be taken into account ($n_{coll} \propto 3 - 5$ would be a typical value for Au-Au collisions at the top RHIC energy).




Figure 4. (Color online) Variation of $v_4/(v_2)^2$ with the Knudsen number.

Qualitatively, in the limit of small n_{coll} (far from equilibrium), one expects both v_2 and v_4 to scale like n_{coll} , so that $v_4/(v_2)^2$ scales like $1/n_{coll}$: we thus expect that the farther the system from equilibrium, the larger $v_4/(v_2)^2$ [16]. In order to have a more quantitative estimate of the effects of partial thermalization, we use a 2 + 1-dimensional solution of the relativistic Boltzmann equation to study systems with arbitrary n_{coll} . We

use the Knudsen number [16], $K \propto 1/n_{coll}$, as a measure of the degree of thermalization of the system.

Figure 4 displays the dependence of $v_4/(v_2)^2$ considering the degree of thermalization of the medium. In the limit $K \rightarrow 0$, transport results show that $v_4/(v_2)^2 = 0.52$, which is close to $1/2$. We also observe, as expected from the low n_{coll} limit, that increasing K leads to an increase of $v_4/(v_2)^2$. But this effect is only a small correction (typically 10% for a central Au-Au collision).




Figure 5. Same plot as figure 2, with one additional curve showing the effect of the deviation from local equilibrium.

The effects of partial thermalization on the centrality dependence of $v_4/(v_2)^2$ are displayed on figure 5. The values of the Knudsen number needed for this plot are taken from a previous study [15]. The value of K is 0.3 for a central Au-Au collision, and increases for more peripheral collisions. Figure 5 shows that adding the effects of deviation from local thermal equilibrium to the fluctuations, our prediction overshoots slightly the data for midcentral and peripheral collisions, but the overall agreement is good. We do not yet understand the large value of $v_4/(v_2)^2$ for central collisions.

7. Conclusion

To conclude, I would like to recall three points: 1) v_4 is mainly induced by v_2 ; 2) the deviation from local equilibrium has a small effect on $v_4/(v_2)^2$; 3) eccentricity fluctuations explain the observed values of $v_4/(v_2)^2$, except for the most central collisions which require further investigation.

Acknowledgments

This work is funded by ‘Agence Nationale de la Recherche’ under grant ANR-08-BLAN-0093-01.

- [1] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **75** (2007) 054906.
- [2] S. Huang [PHENIX Collaboration], J. Phys. G **35** (2008) 104105.
- [3] N. Borghini and J. Y. Ollitrault, Phys. Lett. B **642** (2006) 227
- [4] B. Alver *et al.*, Phys. Rev. C **77** (2008) 014906.
- [5] M. Miller and R. Snellings, arXiv:nucl-ex/0312008.
- [6] R. S. Bhalerao, N. Borghini and J. Y. Ollitrault, Nucl. Phys. A **727** (2003) 373
- [7] J. Y. Ollitrault, A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **80** (2009) 014904
- [8] <http://projects.hepforge.org/tglaubermc/>.
- [9] B. B. Back *et al* [PHOBOS Collaboration], Phys. Rev. C **70** (2004) 021902
- [10] Y. Bai, “Anisotropic Flow Measurements in STAR at the Relativistic Heavy Ion Collider”, PhD thesis, NIKHEF and Utrecht University, 2007.
- [11] R. Lacey, private communication.
- [12] C. Gombeaud and J. Y. Ollitrault, Phys. Rev. C **81**, (2010) 014901
- [13] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **77** (2008) 054901
- [14] J. Adams *et al.* [STAR Collaboration], Phys. Rev. C **72** (2005) 014904
- [15] H. J. Drescher, A. Dumitru, C. Gombeaud and J. Y. Ollitrault, Phys. Rev. C **76** (2007) 024905.
- [16] R. S. Bhalerao, J. P. Blaizot, N. Borghini and J. Y. Ollitrault, Phys. Lett. B **627** (2005) 49.